Table 55 (page 1 of 2). Acquired immunodeficiency syndrome (AIDS) cases, according to race, Hispanic origin, sex, and transmission category for persons 13 years of age and over at diagnosis: United States, selected years 1985–2001

[Data are based on reporting by State health departments]

Race, Hispanic origin, sex, and transmission category	All years¹	All years <sup>1</sup>	1985	1990	1995	1998	1999	2000	January– June 2001	
Race and Hispanic origin	Percent distribution <sup>2</sup>	Number, by year of report								
All races	100.0	758,434	8,028	40,740	69,774	45,514	44,446	40,230	19,002	
Men who have sex with men Injecting drug use	47.1	357,583	5,348	23,658	30,944	16,878	15,632	13,648	6,241	
	24.3	184,247	1,390	9,270	18,802	10,691	9,878	8,099	3,169	
injecting drug use	6.3	48,132	661	2,943	4,185	2,224	1,929	1,587	657	
	0.7	5,171	71	347	467	177	156	98	48	
	10.5	79,769	151	2,253	8,479	7,124	7,344	6,562	2,757	
	3.7	28,368	107	1,484	2,794	1,894	1,769	1,490	611	
	1.1	8,698	165	770	572	277	268	297	96	
	9.9	74,834	242	1,499	6,325	8,143	9,239	9,939	6,034	
White, non-Hispanic	100.0	335,375	4,892	22,062	29,108	15,830	14,622	13,242	6,168	
Men who have sex with men Injecting drug use	67.6	226,714	3,976	16,474	18,826	9,202	8,209	7,239	3,277	
	12.1	40,591	246	2,054	4,145	2,385	2,162	1,852	799	
injecting drug use	7.6	25,427	411	1,646	2,125	1,046	974	765	316	
	1.2	3,939	59	279	329	105	113	73	38	
	5.2	17,445	34	650	1,940	1,332	1,331	1,121	477	
	2.0	6,675	19	349	692	402	401	354	142	
	1.5	5,034	125	505	270	104	85	95	38	
	4.8	16,225	41	454	1,473	1,656	1,748	2,097	1,223	
Black, non-Hispanic	100.0	296,214	1,989	12,796	28,463	21,302	21,589	19,608	9,505	
Men who have sex with men Injecting drug use	27.2	80,555	783	4,453	7,454	4,796	4,676	4,067	1,881	
	35.3	104,468	743	5,170	10,665	6,239	5,714	4,626	1,835	
injecting drug use	5.5	16,243	165	941	1,496	846	688	585	251	
	0.2	685	5	34	81	37	27	14	8	
	16.2	47,965	91	1,216	4,861	4,482	4,717	4,329	1,869	
	5.5	16,348	65	851	1,541	1,137	1,045	910	384	
	0.8	2,436	29	160	205	120	126	148	46	
	14.8	43,862	173	822	3,701	4,782	5,641	5,839	3,615	
Hispanic <sup>6</sup>	100.0	118,012	1,087	5,476	11,365	7,780	7,671	6,764	3,026	
Men who have sex with men Injecting drug use Men who have sex with men and	38.4	45,316	546	2,449	4,182	2,612	2,502	2,093	970	
	32.4	38,185	394	2,010	3,878	1,989	1,938	1,552	506	
injecting drug use	5.0	5,930	83	331	512	301	246	215	70	
	0.4	434	7	28	49	27	13	8	2	
	11.5	13,524	26	375	1,593	1,241	1,215	1,007	376	
	4.3	5,106	23	279	539	336	308	210	78	
	0.8	978	6	82	78	43	46	45	7	
	11.6	13,645	25	201	1,073	1,567	1,711	1,844	1,095	

See footnotes at end of table.

Health, United States, 2002

## Table 55 (page 2 of 2). Acquired immunodeficiency syndrome (AIDS) cases, according to race, Hispanic origin, sex, and transmission category for persons 13 years of age and over at diagnosis: United States, selected years 1985-2001

[Data are based on reporting by State health departments]

Race, Hispanic origin, sex, and transmission category	All years¹	All years <sup>1</sup>	1985	1990	1995	1998	1999	2000	January– June 2001		
Sex	Percent distribution <sup>2</sup>	Number, by year of report									
Male	100.0	629,429	7,504	36,193	56,776	35,104	34,094	30,251	14,304		
Men who have sex with men Injecting drug use	56.8 21.0	357,583 132,238	5,348 1,103	23,658 6,923	30,944 13,376	16,878 7,440	15,632 6,893	13,648 5,554	6,241 2,215		
injecting drug use	7.6 0.8 4.5 1.4 0.8 8.5	48,132 4,893 28,430 8,931 4,944 53,209	661 68 32 25 102 190	2,943 332 715 454 440 1,182	4,185 438 2,924 871 319 4,590	2,224 153 2,723 645 151 5,535	1,929 143 2,947 634 137 6,413	1,587 93 2,537 514 146 6,686	657 45 1,077 228 54 4,015		
Female	100.0	129,005	524	4,547	12,998	10,410	10,352	9,979	4,698		
Injecting drug use	40.3 0.2 39.8 15.1 2.9 16.8	52,009 278 51,339 19,437 3,754 21,625	287 3 119 82 63 52	2,347 15 1,538 1,030 330 317	5,426 29 5,555 1,923 253 1,735	3,251 24 4,401 1,249 126 2,608	2,985 13 4,397 1,135 131 2,826	2,545 5 4,025 976 151 3,253	954 3 1,680 383 42 2,019		

<sup>&</sup>lt;sup>1</sup>Includes cases prior to 1985 and through June 30, 2001.

<sup>6</sup>Persons of Hispanic origin may be of any race.

NOTES: The AIDS case reporting definitions were expanded in 1985, 1987, and 1993. See Appendix II, AIDS. Excludes data for U.S. dependencies and possessions and independent nations in free association with the United States. Data for all years have been updated through June 30, 2001, to include temporally delayed case reports and may differ from previous editions of Health, United States. Similar data as of December 31, 2001, are available in the Centers for Disease Control and Prevention, HIV/AIDS Surveillance Report, Year-end edition Vol 13 No 2. 2001.

SOURCE: Centers for Disease Control and Prevention, National Center for HIV, STD, and TB Prevention, Division of HIV/AIDS Prevention—Surveillance and Epidemiology, AIDS Surveillance, 2001 special data run.

184 Health, United States, 2002

<sup>&</sup>lt;sup>2</sup>Percents may not sum to 100 percent due to rounding.
<sup>3</sup>Includes persons who have had heterosexual contact with a person with human immunodeficiency virus (HIV) infection or at risk of HIV infection.

<sup>&</sup>lt;sup>4</sup>Receipt of blood transfusion, blood components, or tissue.

<sup>&</sup>lt;sup>5</sup>Includes persons for whom risk information is incomplete (because of death, refusal to be interviewed, or loss to followup), persons still under investigation, men reported to have had heterosexual contact only with prostitutes, and interviewed persons for whom no specific risk is identified.